

RESVERATROL ATTENUATES EXERCISE-INDUCED ADAPTIVE RESPONSES IN RATS SELECTIVELY BRED FOR LOW RUNNING PERFORMANCE

Nikolett Hart¹, Linda Sarga¹, Zsolt Csende¹, Lauren G. Koch², Steven L. Britton², Kelvin J.A. Davies³, Zsolt Radak¹ □ ¹Research Institute of Sport Science, Semmelweis University, Budapest, Hungary, ²Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, Michigan, USA, ³Ethel Percy Andrus Gerontology Center of the Davis School of Gerontology; and Division of Molecular & Computational Biology, Department of Biological Sciences of the College of Letters, Arts & Sciences: the University of Southern California, Los Angeles, USA

□ Low capacity runner (LCR) rats have been developed by divergent artificial selection for treadmill endurance capacity to explore an aerobic biology-disease connection. The beneficial effects of resveratrol supplementation have been demonstrated in endurance running. In this study it was examined whether 12 weeks of treadmill exercise training and/or resveratrol can retrieve the low running performance of the LCR and impact mitochondrial biogenesis and quality control. Resveratrol regressed running performance in trained LCR ($p < 0.05$). Surprisingly, exercise and resveratrol treatments significantly decreased pAMPK/AMPK, SIRT1, SIRT4, forkhead transcription factor 1 (FOXO1) and mitochondrial transcription factor A (TFAM) levels in these animals ($p < 0.05$). Mitochondrial fusion protein, HSP78 and polynucleotide phosphorylase were significantly induced in LCR-trained, LCR-resveratrol treated, LCR-trained and resveratrol treated groups compared to LCR-controls. The data indicate that the AMPK-SIRT1-NAMPT-FOXO1 axis could be important to the limited aerobic endurance capacity of low running capacity rats. Resveratrol supplementation was not beneficial in terms of aerobic endurance performance, mitochondrial biogenesis, or quality control.

Key words: Resveratrol, Exercise, Sirtuins, Adaptation, Free Radicals

INTRODUCTION

Large-scale epidemiologic studies demonstrate a strong statistical association between low capacity for aerobic exercise and increased risk for development of complex diseases (DeMarco *et al.* 2012). Although an underlying mechanistic relationship has been hypothesized (aerobic hypothesis), the complexity of both aerobic metabolism and the putative disease conditions in humans makes the unraveling of cause and effect a challenge (Kivela *et al.* 2010). In 1996 Koch and Britton (Koch and Britton 2001) initiated a prospective test of the linkage between aerobic capacity and disease risk by applying large-scale artificial selective breed-

Address correspondence to Zsolt Radak, Ph.D, Institute of Sport Science, Faculty of Physical Education and Sport Science, Semmelweis University, Alkotás u. 44. TF Budapest, Hungary; Tel: +36 1 3565764; Fax: +36 1 356 6337; Email: radak@tf.hu

ing in rats with widely varying genetic backgrounds to produce low and high strains that differ for intrinsic (i.e., untrained) aerobic endurance treadmill running capacity (Koch and Britton 2001). The hypothesis was that rats selectively bred as Low Capacity Runners (LCR) would display disease risks and rats bred as High Capacity Runners (HCR) would have positive health effects. Several studies report that the LCR present with negative health features including metabolic syndrome, reduced heart function, hepatic steatosis, disordered sleep, and diminished oxidative capacity in skeletal muscle (Wisloff *et al.* 2005; Thyfault *et al.* 2009; Kivela *et al.* 2010). In contrast, HCR demonstrate greater maximal oxygen consumption, insulin sensitivity and improved metabolic health. In accordance with the aerobic hypothesis, aged rats with low intrinsic aerobic capacity have diminished longevity and display a reduced ability for mitochondrial regeneration, decreased metabolic control in the heart, and reduced antioxidant status (Koch and Britton 2001). Therefore, this model is excellent to study how life-style interventions, such as exercise training or nutritional manipulation, could overcome or exacerbate the effects of genetics on physiological and biochemical processes that lead to health promotion (Kamei *et al.* 2004; Wisloff *et al.* 2005; Bowman *et al.* 2010).

Exercise training and resveratrol have been suggested for therapeutic potential in treatment of metabolic disorders. Both interventions increase oxidative metabolism in skeletal muscle by induction of a highly integrated molecular network which results in increased insulin sensitivity (Nunn *et al.* 2010), activity of AMP-activated protein kinase, (AMPK) peroxisome proliferator-activated receptor- γ coactivator 1 α (PGC-1 α), and mitochondrial content (Baur *et al.* 2006; Li *et al.* 2011). Recent findings reveal that knockdown of a mitochondrial located sirtuin, SIRT4, results in increased fatty acid oxidation, enhanced mitochondrial function, and higher AMPK levels in skeletal muscle (Nasrin *et al.* 2010), suggesting a primary role of this sirtuin in aerobic metabolism. Moreover, another member of the sirtuin family, the NAD⁺ dependent SIRT1, is also an important regulator of oxidative mitochondrial metabolism (Rodgers *et al.* 2008) by deacetylation of the peroxisome proliferator-activated receptor gamma co-activator 1 alpha (PGC-1 α) (Nemoto *et al.* 2005). PGC-1 α activates gene transcription factors important for mitochondrial biogenesis including nuclear respiratory factor 1 (NRF1) and mitochondrial transcription factor A (TFAM), which encode nuclear and mitochondrial DNA imprinted genes required for the production of mitochondrial proteins. Mitochondrial fusion and fission are important mechanisms for maintenance of the mitochondrial network and for quality control (Westermann 2010), and thus impact mitochondrial function (Otera and Mihara 2011). The quality control of mitochondrial proteins is supervised by Lon protease and HSP78, which prevent the accumula-

tion of oxidized and dysfunctional proteins in mitochondria (Bota and Davies 2002; Rottgers *et al.* 2002; Ngo and Davies 2009). Recent studies have revealed that polynucleotide phosphorylase (PNPase) is involved in the stability of mitochondria (Chen *et al.* 2006) and is crucial for the import of nuclear coded RNAs into the mitochondrial matrix (Wang *et al.* 2010).

It is well known that similar exercise programs can result in different training responses for different subjects (Radak *et al.* 2013). However, the possible different responses to nutritional and pharmacological interventions, including resveratrol treatment, remain to be well characterized. Resveratrol has been shown to reduce plasma triglyceride concentrations, oxidative stress, and inflammation in humans (Zern *et al.* 2005), as well as to improve insulin sensitivity in rats (Zheng *et al.* 2012).

Therefore, in the present study, the LCR contrasting rat model system was utilized to test the influence of exercise training, and resveratrol singularly, and the combined effects of training and resveratrol, upon indices of health, including running performance, VO_2max , and forearm gripping strength. We were particularly interested to further understand retrieval of the negative LCR clinical phenotype. Positive effects of exercise training and resveratrol treatment on the LCR group would mean that life-style interventions could compensate, in some degree, for the health treating effects of a poor genetic setup. Both physical exercise and resveratrol can modulate the generation and activity of reactive oxygen species (ROS), which are implicated in a variety of diseases, and signaling, and mitochondrial biogenesis (Radak *et al.* 2013). Therefore, as an additional aim, the relative density of mtDNA and activities of a number of factors that regulate mitochondrial biogenesis, quantity and quality control and differentiation from gastrocnemius muscle, were investigated for possible explanations.

METHODS

Animals

Artificial selective breeding, starting with a founder population of 186 genetically heterogeneous rats (N:NIH stock), was used to develop rat strains differing in inherent aerobic capacity. The procedure has been described in detail previously (Koch and Britton 2001). Briefly, endurance running capacity was assessed on a treadmill and the total distance run during a speed-ramped exercise test was used as a measure of maximal aerobic capacity. Rats with the lowest capacity were bred to produce the LCR strain. A subgroup of 24 male rats from generation 22 was phenotyped for intrinsic treadmill running capacity when 11 weeks old, at the University of Michigan (Ann Arbor, USA), and then shipped via air freight to Semmelweis University (Budapest, Hungary) for further study.

Investigations were carried out according to the requirements of The Guiding Principles for Care and Use of Animals, EU, and approved by the ethics committee of Semmelweis University.

Exercise protocol and resveratrol treatment

Twenty four LCR male rats, aged 13 months, were equally assigned to control (LCR-C), trained (LCR-Tr), resveratrol treated control (LCR-Rsv), and trained resveratrol treated (LCR-TrRsv), groups. Control rats had access to a treadmill with an electrical stimulator, three times a week, for ten min. Trained rats were introduced to treadmill running for three days, then for the next two weeks the running speed was set to 10 m/min, on a 5% incline for 30 min. The treadmill was equipped with a high pressure air pipe and electric grid to stimulate running.

In the following week, maximal oxygen uptake ($VO_2\text{max}$) was measured on a motor driven treadmill (Columbus Inst. Columbus, Ohio) with a gradually increasing intensity. $VO_2\text{max}$ was measured for each animal, using three criteria: (i) no change in VO_2 when speed was increased, (ii) rats no longer kept their position on the treadmill, and (iii) respiratory quotient ($RQ = VCO_2/VO_2$) > 1. Based on the level of $VO_2\text{max}$, a treadmill speed corresponding to 60% $VO_2\text{max}$ was determined and used for daily training for one hour, five times per week. $VO_2\text{max}$ was measured every second week and running speed was adjusted accordingly. The total training period lasted 12 weeks. In addition, the forelimb strength of the animals was assessed weekly by using a gripping test as described by Marton *et al.* (Marton *et al.* 2010). This test measures the length of time the animals are able to hang on a pole.

Resveratrol supplementation (100 mg/kg, oral dose) (Smith *et al.* 2009) was started two weeks before habitual treadmill running was introduced to the animals, and four weeks before the actual training started, thereby lasting 16 weeks. The animals were sacrificed two days after the last exercise session to avoid the acute metabolic effects of the final run. The gastrocnemius muscle was dissected and homogenized in buffer (HB) containing 137 mM NaCl, 20mM Tris-HCl (pH 8.0), 2% NP 40, 10% glycerol and protease inhibitors.

ROS, and protein carbonyl content and antioxidant enzyme activities

Intracellular oxidant and redox-active iron levels (Kalyanaraman *et al.* 2012) were estimated using modifications of the dichlorodihydrofluorescein diacetate (H_2DCFDA) staining method (Radak *et al.* 2004). The oxidative conversion of stable, nonfluorometric DCF-DA to highly fluorescent 2'7'-dichlorofluorescein (DCF) was measured in the presence of esterases, as previously reported (Radak *et al.* 2004). This assay approximates levels of reactive species, such as superoxide radical, hydroxyl radical, and hydrogen peroxide. The method has been widely used in the lit-

erature but does have the problem of not being particularly specific, and results can be strongly affected by release of labile iron or copper (Kalyanaraman *et al.* 2012). Briefly, the H₂DCFDA (Invitrogen-Molecular Probes #D399) was dissolved to a concentration of 12.5 mM in ethanol and kept at -80 °C in the dark. The solution was freshly diluted with potassium phosphate buffer to 125 μM before use. For fluorescence reactions, 96-well, black microplates were loaded with potassium phosphate buffer (pH 7.4) to a final concentration of 152 μM/well. Then eight μl diluted tissue homogenates and 40 μl 125 μM dye were added to achieve a final dye concentration of 25 μM. The change in fluorescence intensity was monitored every five minutes for 30 minutes with excitation and emission wavelengths set at 485 nm and 538 nm (Fluoroskan Ascent FL) respectively. Data obtained after 15 min were used. The fluorescence intensity unit was normalized with the protein content and expressed in relative unit production per minute. The protein carbonyl measurement was assessed as described earlier (Koltai *et al.* 2012).

Western blots

Ten to 50 micrograms of protein were electrophoresed on 8-12% v/v polyacrylamide SDS-PAGE gels. Proteins were electrotransferred onto PVDF membranes. The membranes were subsequently blocked and incubated at room temperature with antibodies (1:500 #sc-13067 Santa Cruz PGC-1 (H-300), 1:1400 #2532 Cell Signaling AMPKα, 1:500 #2535 Cell Signaling p-AMPKα (Thr172) (40H9), 1:1000 #sc-33771 Santa Cruz NRF-1 (H-300), 1:500 #sc-30963 Santa Cruz mtTFA (E-16) /TFAM/, 1:500 #sc-98900 Santa Cruz Fis1 (F1-152), 1:10000 #sc-50330 Santa Cruz Mfn1 (H-65), 1:1000 #sc-99006 Santa Cruz PNPase (H-124), 1:200 #U7757 Sigma-Aldrich UCP3, 1:5000 #ab87253 Abcam CLPB /HSP78/, 1: 15000, #T6199 Sigma alpha-tubulin). The antibody for Lon protease was generated in our laboratory, as described previously (Merrill *et al.* 1997). After incubation with primary antibodies, membranes were washed in TBS-Tween-20 and incubated with HRP-conjugated secondary antibodies. After incubation with the secondary antibody, membranes were repeatedly washed. Membranes were incubated with an ECL Plus reagent (RPN 2132, Amersham) and protein bands were visualized on X-ray films. The bands were quantified by ImageJ software, and normalized to tubulin, which served as an internal control.

Assessment of SIRT1 activity

To measure SIRT1 deacetylase activity, a Cyclex SIRT1/Sir2 Deacetylase Fluorometric Assay Kit (Cyclex, CY-1151) was used according to the established protocol including the separation of nuclear extract (Koltai *et al.* 2012). The purity of nuclear extract was checked by organelle specific antibodies as reported earlier (Radak *et al.* 2009).

To prepare nuclear fractions, the homogenate was centrifuged at $1000 \times g$ for 10 min at 4 °C, and the pellet was suspended in HB and re-centrifuged. The pellet was then re-suspended in HB with 0.5% NP40 and again centrifuged. Next, the pellet was washed twice in HB. After centrifugation, the final nuclear pellet was rocked for 30 min after the addition of a 1/10 (vol/vol) of 2.5 M KCl and centrifuged at 14,000 rpm for 30 min. For the measurement of SIRT1, five microliters of nuclear extracts of rat gastrocnemius muscle were mixed with a reaction mixture (40 μ l) containing 50 mM Tris-HCl pH 8.8, 4 mM MgCl₂, 0.5 mM DTT, 0.25 mAU/ml Lysyl endopeptidase, 1 μ M Trichostatin A, 20 μ M Fluoro-Substrate Peptide, and 200 μ M NAD⁺ on a microplate. The samples were mixed and incubated for ten min at room temperature and the fluorescence intensity (ex. 355nm, em. 460 nm) was read every ten min for two hours and normalized by the protein content (Koltai *et al.* 2010).

Measurement of mtDNA by PCR

The mtDNA content was quantified as the mtDNA to nuclear DNA (nDNA) ratio (mtDNA/nDNA). Total DNA was extracted (Fast DNA kit #6540-400 BIO 101 Systems Qiogene) and quantified spectrophotometrically. The mtDNA content was measured by PCR (Rotor-Gene 6000, Corbett Research, Australia) using the following conditions: 94 °C for two min (initial denaturation), 94 °C for 30 s, 60 °C for 30 s, 72 °C for 45 s (25 cycles), 72 °C for ten min (final extension), and corrected by the simultaneous measurement of a single copy nuclear BDNF gene. Primers used for the analysis of mtDNA were R-CYTB-F (5'-CCC CAG AGG ATT AAA CTC CAA CGC A-3'), and R-CYTB-R (5'-GGG TGG GGT CAG GGG GT-3'). Primers used for the analysis of nDNA were R-BDNF-genom-exon-IV-F (5'-TTG GGA TGG GAA AGA TGG G-3'), and R-BDNF-genom-exon-IV-R (5'-CAG AGT AGG AGG GAA CAA GTG TGA C-3'). The mtDNA content was normalised to nDNA. Data are expressed as the mean of three measurements.

Statistical analyses

Because of the limited sample size and for the purpose of finding the appropriate statistical procedure, normality was tested on all dependent variables (Shapiro Wilk's W-test). It was found that few of the dependent variables demonstrated a normal distribution, namely: NRF1, Fis1, NAMPT, LonP, Mfn1, and VO₂max. Therefore, non-parametric Kruskal-Wallis ANOVA analysis was used to test for differences among the dependent variables. The Mann-Whitney U-test was applied for post-hoc analyses. Significance level was set at $p < 0.05$.

RESULTS

Exercise performance

The body mass of the LCR-Tr group was significantly lower ($p < 0.001$) ($470 \pm 47\text{g}$) than LCR-C ($609 \pm 38\text{g}$).

Exercise training significantly increased the level of running distance of LCR-Tr ($p < 0.01$) compared to control and resveratrol treated rats ($p < 0.05$) (Fig. 1A) and similar results were present for the levels of VO_2max (Fig. 1B). The gripping strength of the upper limbs did not show significant differences between groups (Fig. 1C).

Effects of exercise and resveratrol on mitochondrial metabolic factors

AMPK activity is a known modulator of PGC-1 α . Surprisingly, exercise and resveratrol treatments significantly decreased pAMPK/AMPK levels in LCR rats compared to control animals ($p < 0.05$) (Fig. 2A). Interestingly, the changes in SIRT1 levels were very similar in pattern to AMPK activity although to a lesser degree (Fig. 2B). Nicotinamide phos-




FIG. 1. Running distance (A), VO_2max (B), and gripping time (C) of LCR rats. Running distance (m) (A), maximal oxygen uptake (VO_2max ; ml/kg/min) (B) and gripping time (s) (C) for low capacity runner (LCR) rats were measured every second week across a 12 week exercise training period and during the two weeks of treadmill habituation. Control LCR (LCR-C), trained LCR (LCR-Tr), resveratrol treated control LCR (LCR-Rsv), trained resveratrol treated LCR (LCR-TrRsv) groups. Values are means \pm SD for six animals per group *Significantly different from control group, + Significantly different from trained group, # Significantly different from resveratrol treated group, $p < 0.05$.




FIG. 2. The effect of exercise and resveratrol on metabolic factors in skeletal muscle of LCR rats. The activity of AMPK, assessed by the pAMPK/AMPK ratio (A), which decreased with training and resveratrol treatment. The activity of SIRT1 was not changed by exercise or resveratrol (B). The protein levels of NAMPT (C), PGC-1 α (D), FOXO1 (E) and SIRT4 (F) were determined by Western blot and densitometry in gastrocnemius muscle. G panel shows the mtDNA content. Control LCR (LCR-C), trained LCR (LCR-Tr), resveratrol treated control LCR (LCR-Rsv), trained resveratrol treated LCR (LCR-TrRsv) groups. Values are means \pm SD for six animals per group *Significantly different from control group, + Significantly different from trained group, # Significantly different from resveratrol treated group, $p < 0.05$.

phosphoryltransferase (NAMPT) was measured because it is involved in the biosynthesis of NAD⁺ and SIRT1 is dependent on the availability of NAD⁺. The NAMPT levels were decreased in both exercising groups ($p < 0.05$) (Fig. 2C). Despite the decreasing activity of SIRT1, PGC-1 α levels increased with exercise training (Fig. 2D). FOXO1 is an important transcription factor for regulation of gluconeogenesis and adipogenesis and in this model resveratrol alone and with training decreased FOXO1 levels (Fig 2E). Mitochondrial sirtuin, SIRT4, is related to fat metabolism and the data revealed that exercise training and resveratrol treatment decreased SIRT4 content (Fig 2F). mtDNA levels increased significantly with exercise and resveratrol (Fig 2G).

The effects of exercise and resveratrol on mitochondrial quantity and quality control

TFAM protein levels decreased in LCR-Tr and LCR-TrRsv rats compared to controls ($p < 0.05$) (Fig. 3A). The concentration of NRF-1 was not affected by exercise training or resveratrol treatments (Fig. 3B). Mfn1 is




FIG. 3. Markers of quantity and quality control of mitochondria. Exercise and resveratrol altered the levels of TFAM (A) while the NRF1 concentration was unchanged (B). We evaluated mitochondrial fusion, Mfn1(C), and fission Fis1 (D) and quality control was assessed by Lon protease (E), HSP78 (F) and PNPase (G) levels. Control LCR (LCR-C), trained LCR (LCR-Tr), resveratrol treated control LCR (LCR-Rsv), trained resveratrol treated LCR (LCR-TrRsv) groups. Values are means \pm SD for six animals per group *Significantly different from control group, + Significantly different from trained group, # Significantly different from resveratrol treated group, $p < 0.05$.

a mediator of mitochondrial fusion and was significantly induced in LCR-Tr, LCR-Rsv, and LCR-TrRsv groups compared to LCR-C (Fig. 3C). The mitochondrial fission controlling protein, Fis1, was not affected by exercise training in any of the LCR groups (Fig 3D). Lon protease was measured to assess the level of protein degradation in mitochondria, which could be important for quality control. However, significant differences were not noted in all groups (Fig. 3E). HSP78 and PNPase were induced by both training and resveratrol (Fig. 3F, Fig. 3G).

Oxidative stress markers

The levels of ROS, as measured by the fluorescent activity of H_2DCFDA , increased with exercise training (Fig. 4A). The levels of carbonyl groups were assessed as potential markers of oxidative protein damage and the data clearly indicate decreased levels in LCR-TrRsv rats compared to controls ($p < 0.05$) (Fig. 4B).

DISCUSSION

Low running capacity rats have shown a wide range of physiological disorders (Wisloff *et al.* 2005; Thyfault *et al.* 2009; Kivela *et al.* 2010), which




FIG. 4. Oxidative stress markers. Rat gastrocnemius muscle was stained with dichlorodihydrofluorescein diacetate (H_2DCFDA) to measure relative steady-state oxidant levels and redox-active iron release levels (both increase DCF fluorescence) as an estimate of levels of reactive oxygen species (ROS) (A). The oxidative damage of proteins was evaluated by protein carbonyl groups (B). Control LCR (LCR-C), trained LCR (LCR-Tr), resveratrol treated control LCR (LCR-Rsv), trained resveratrol treated LCR (LCR-TrRsv) groups. Values are means \pm SD for six animals per group *Significantly different from control group, + Significantly different from trained group, # Significantly different from resveratrol treated group, $p < 0.05$.

suggests that the increases in aerobic performance could be important to overcome some of these disorders. We have demonstrated in this study, that exercise training alone can increase VO_2 max of LCR rats. Since higher VO_2 max is associated with longer life-spans of LCR and HCR rats (Koch and Britton 2001) it can be suggested that the enhanced endurance ability would result in an increased life-span for LCR rats and reduce the metabolic disorders associated with the low endurance capacity of LCR rats.

In the present model resveratrol supplementation was not beneficial, although according to the general view, resveratrol supplementation promotes health. However, it must be noted that in some models resveratrol supplementation has been reported to act differently on obese prone and obese resistant rats (Louis *et al.* 2012). Therefore, the response to resveratrol treatment could be highly selective and could be based on metabolic status.

One of the striking observations of this study was that exercise training decreased AMPK activity in this rat model. AMPK acts as an energy sensor to cope with metabolic challenges and can interfere with fuel preference and availability (Kurth-Kraczek *et al.* 1999). Therefore, the exer-

cise related decreased activity of AMPK could, at least in part, explain the limited running capacity of LCR rats.

It has been shown that exercise training increases NAMPT levels in skeletal muscle of wild rats (Koltai *et al.* 2010) and humans (Costford *et al.* 2010). NAMPT is important to NAD production which serves as a co-factor for sirtuins. Indeed, the activity of SIRT1 tends to decrease in a similar pattern to NAMPT and AMPK. AMPK has been shown to be linked to the activity of SIRT1, since AMPK can activate SIRT1 by increasing the levels of NAD⁺ (Canto *et al.* 2009) which, as a metabolic sensor, can control the expression of a number of genes involved in the metabolism of skeletal muscle (Canto *et al.* 2009). The AMPK associated decreased activity of SIRT1 could lead to increased acetylation of FOXO1, which is a target for SIRT1 deacetylase (Canto *et al.* 2009). Increased acetylation of FOXO1 attenuates the activity of this transcription factor (Nakae *et al.* 2012). The present data show decreased activity of AMPK and SIRT1 and decreased content of NAMPT and FOXO1 with exercise training. We strongly believe that this “unusual” response to exercise causes the limitation of aerobic capacity which limits the endurance capacity of low running capacity rats. This suggestion is based on the fact that wild rats respond to exercise training with increased AMPK-SIRT1-NAMPT (Costford *et al.* 2010; Koltai *et al.* 2010) and FOXO1 levels (Canto *et al.* 2009). In addition, other data from our laboratories on HCR rats suggests up-regulation of the AMPK-SIRT1-NAMPT-FOXO1 system (Hart *et al.* 2013).

The exact function of mitochondrial located SIRT4 is still in debate, but a recent study revealed that knockdown of SIRT4 by shRNA significantly increased fat metabolism in myotubes (Nasrin *et al.* 2010). This is an important finding and could indicate that lower levels of SIRT4 result in increased free fatty acid utilization, a key element for increased endurance capacity (Holloszy and Coyle 1984; Spina *et al.* 1996). Our present observations in trained and resveratrol supplemented rats suggests that the high body mass observed in control LCR rats could be due to impaired utilization of fat, which is enhanced by exercise training and resveratrol supplementation. Moreover, it is also known that SIRT4 is involved in the development of insulin resistance (Chen *et al.* 2010). Therefore, the down-regulating effects of exercise could be meaningful, especially in LCR rats, which suffer from this problem.

TFAM levels and the activity of SIRT1 changed in similar patterns with resveratrol and exercise intervention, suggesting a functional relationship between these proteins. TFAM is reported to increase the half life of and stabilize mtDNA, and thus increased levels of TFAM could have beneficial effects (Ikeuchi *et al.* 2005). However, the decreased levels of TFAM and mtDNA content with training, indicate an increased vulnerability of mtDNA.

Mitochondrial fission and fusion interact with each other. Fusion enables mitochondria to mix their contents within interconnected mitochondrial reticulum in order to minimize abnormalities. Fusion pathways have been shown to have beneficial roles in muscular atrophy (Romanello *et al.* 2010), endurance (Garnier *et al.* 2005), lifespan (Scheckhuber *et al.* 2007) and ROS production (Yu *et al.* 2006). In the present study, we have found that both exercise training and resveratrol enhance the levels of the main fusion protein, indicating beneficial affects of these interventions on the mitochondrial network.

Lon protease plays an important role in the quality control of mitochondrial proteins by degrading oxidatively modified proteins, and HSP78, an important mitochondria chaperone involved in LonP- associated degradation and quality control (Rottgers *et al.* 2002), and appears to be very well induced by exercise and resveratrol treatment in these LCR rats. Indeed, it has been suggested that long-term health conditions is accomplished by a complex network called vitagenes, a group of genes involved in preserving cellular homeostasis during stressful conditions (Calabrese *et al.* 2011; Calabrese *et al.* 2012). Vitagenes encode for heat shock proteins, sirtuins and Lon protease as well (Calabrese *et al.* 2006; Cornelius *et al.* 2013), and our currenet findings suggest an improved level of quality control in the mitochondria with resveratrol and exercise training.

PNPase content was evaluated because it is important for the homeostasis of mitochondria (Chen *et al.* 2006) and can induce cellular senescence (Sarkar *et al.* 2005). Ablation of PNPase leads to a significant drop in enzymatic activities of respiratory complexes and decreases ATP production, leading to activation of AMPK phosphorylation (Chen *et al.* 2006). PNPase knockdown results in impaired mitochondrial membrane potential, which can readily affect fusion and fission (Chen *et al.* 2006). In the present study it was observed that exercise training increased PNPase content, suggesting that exercise played a role in the improvement of mitochondrial respiration.

In summary, the present data show that exercise intervention is an excellent tool to overcome genetic weaknesses by increasing endurance capacity. Exercise was able to overcome some of the weaknesses of mitochondrial systems through the activation of mitochondrial biogenesis. However, our data revealed that the AMPK-SIRT1-NAMPT-FOXO1 axis could be important for the limited aerobic endurance capacity of low running capacity rats. Resveratrol supplementation was not beneficial, in terms of aerobic endurance performance, mitochondrial biogenesis, or quality control.

ACKNOWLEDGEMENT

This work was supported by Hungarian grants from ETT 38388, OTKA (K75702) TAMOP-4.2.2/B-10/1-2010-0013 awarded to Z. Radák. The LCR rat model system was supported by grant R24 RR017718 from the National Center for Research Resources of the National Institutes of Health (to L.G.K. and S.L.B.), and National Institutes of Health grant RO1 DK077200 (to S.L.B.). The LCR model can be made available for collaborative study (contact: brittons@umich.edu or lgkoch@umich.edu).

REFERENCES

- Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, Boss O, Gwinn D, Wang M, Ramaswamy S, Fishbein KW, Spencer RG, Lakatta EG, Le Couteur D, Shaw RJ, Navas P, Puigserver P, Ingram DK, de Cabo R and Sinclair DA. 2006. Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 444: 337-342
- Bota DA and Davies KJ. 2002. Lon protease preferentially degrades oxidized mitochondrial aconitase by an ATP-stimulated mechanism. *Nat Cell Biol* 4: 674-680
- Bowman TA, Ramakrishnan SK, Kaw M, Lee SJ, Patel PR, Golla VK, Bourey RE, Haram PM, Koch LG, Britton SL, Wisloff U, Lee AD and Najjar SM. 2010. Caloric restriction reverses hepatic insulin resistance and steatosis in rats with low aerobic capacity. *Endocrinology* 151: 5157-5164
- Calabrese V, Butterfield DA, Scapagnini G, Stella AM and Maines MD. 2006. Redox regulation of heat shock protein expression by signaling involving nitric oxide and carbon monoxide: relevance to brain aging, neurodegenerative disorders, and longevity. *Antioxid Redox Signal* 8: 444-477
- Calabrese V, Cornelius C, Cuzzocrea S, Iavicoli I, Rizzarelli E and Calabrese EJ. 2011. Hormesis, cellular stress response and vitagenes as critical determinants in aging and longevity. *Mol Aspects Med* 32: 279-304
- Calabrese V, Cornelius C, Dinkova-Kostova AT, Iavicoli I, Di Paola R, Koverech A, Cuzzocrea S, Rizzarelli E and Calabrese EJ. 2012. Cellular stress responses, hormetic phytochemicals and vitagenes in aging and longevity. *Biochim Biophys Acta* 1822: 753-783
- Canto C, Gerhart-Hines Z, Feige JN, Lagouge M, Noriega L, Milne JC, Elliott PJ, Puigserver P and Auwerx J. 2009. AMPK regulates energy expenditure by modulating NAD⁺ metabolism and SIRT1 activity. *Nature* 458: 1056-1060
- Chen HW, Rainey RN, Balatoni CE, Dawson DW, Troke JJ, Wasiak S, Hong JS, McBride HM, Koehler CM, Teitell MA and French SW. 2006. Mammalian polynucleotide phosphorylase is an intermembrane space RNase that maintains mitochondrial homeostasis. *Mol Cell Biol* 26: 8475-8487
- Chen YR, Fang SR, Fu YC, Zhou XH, Xu MY and Xu WC. 2010. Calorie restriction on insulin resistance and expression of SIRT1 and SIRT4 in rats. *Biochem Cell Biol* 88: 715-722
- Cornelius C, Crupi R, Calabrese V, Graziano A, Milone P, Pennisi G, Radak Z, Calabrese EJ and Cuzzocrea S. 2013. TRAUMATIC BRAIN INJURY (TBI): OXIDATIVE STRESS AND NEUROPROTECTION. *Antioxid Redox Signal*
- Costford SR, Bajpeyi S, Pasarica M, Albarado DC, Thomas SC, Xie H, Church TS, Jubrias SA, Conley KE and Smith SR. 2010. Skeletal muscle NAMPT is induced by exercise in humans. *Am J Physiol Endocrinol Metab* 298: E117-126
- DeMarco VG, Johnson MS, Ma L, Pulakat L, Mugerfeld I, Hayden MR, Garro M, Knight W, Britton SL, Koch LG and Sowers JR. 2012. Overweight female rats selectively bred for low aerobic capacity exhibit increased myocardial fibrosis and diastolic dysfunction. *Am J Physiol Heart Circ Physiol* 302: H1667-1682
- Garnier A, Fortin D, Zoll J, N'Guessan B, Mettauer B, Lampert E, Veksler V and Ventura-Clapier R. 2005. Coordinated changes in mitochondrial function and biogenesis in healthy and diseased human skeletal muscle. *FASEB J* 19: 43-52
- Hart N, Sarga L, Csende L, Koltai E, Koch LG, Britton SL, Davies KJ, Kouretas D, Wessner B and Radak Z. 2013. Resveratrol enhances exercise training responses in rats selectively bred for high running performance. *Food Chem Toxicol*

- Holloszy JO and Coyle EF. 1984. Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *J Appl Physiol* 56: 831-838
- Ikeuchi M, Matsusaka H, Kang D, Matsushima S, Ide T, Kubota T, Fujiwara T, Hamasaki N, Takeshita A, Sunagawa K and Tsutsui H. 2005. Overexpression of mitochondrial transcription factor α ameliorates mitochondrial deficiencies and cardiac failure after myocardial infarction. *Circulation* 112: 683-690
- Kalyanaraman B, Darley-Usmar V, Davies KJ, Dennery PA, Forman HJ, Grisham MB, Mann GE, Moore K, Roberts LJ, 2nd and Ischiropoulos H. 2012. Measuring reactive oxygen and nitrogen species with fluorescent probes: challenges and limitations. *Free Radic Biol Med* 52: 1-6
- Kamei Y, Miura S, Suzuki M, Kai Y, Mizukami J, Taniguchi T, Mochida K, Hata T, Matsuda J, Aburatani H, Nishino I and Ezaki O. 2004. Skeletal muscle FOXO1 (FKHR) transgenic mice have less skeletal muscle mass, down-regulated Type I (slow twitch/red muscle) fiber genes, and impaired glycemic control. *J Biol Chem* 279: 41114-41123
- Kivela R, Silvennoinen M, Lehti M, Rinnankoski-Tuikka R, Purhonen T, Ketola T, Pullinen K, Vuento M, Mutanen N, Sartor MA, Reunanen H, Koch LG, Britton SL and Kainulainen H. 2010. Gene expression centroids that link with low intrinsic aerobic exercise capacity and complex disease risk. *FASEB J* 24: 4565-4574
- Koch LG and Britton SL. 2001. Artificial selection for intrinsic aerobic endurance running capacity in rats. *Physiol Genomics* 5: 45-52
- Koltai E, Hart N, Taylor AW, Goto S, Ngo JK, Davies KJ and Radak Z. 2012. Age-associated declines in mitochondrial biogenesis and protein quality control factors are minimized by exercise training. *Am J Physiol Regul Integr Comp Physiol* 303: R127-134
- Koltai E, Szabo Z, Atalay M, Boldogh I, Naito H, Goto S, Nyakas C and Radak Z. 2010. Exercise alters SIRT1, SIRT6, NAD and NAMPT levels in skeletal muscle of aged rats. *Mech Ageing Dev* 131: 21-28
- Kurth-Kraczek EJ, Hirshman MF, Goodyear LJ and Winder WW. 1999. 5' AMP-activated protein kinase activation causes GLUT4 translocation in skeletal muscle. *Diabetes* 48: 1667-1671
- Li L, Pan R, Li R, Niemann B, Aurich AC, Chen Y and Rohrbach S. 2011. Mitochondrial biogenesis and peroxisome proliferator-activated receptor- γ coactivator-1 α (PGC-1 α) deacetylation by physical activity: intact adipocytokine signaling is required. *Diabetes* 60: 157-167
- Louis XL, Thandapilly SJ, MohanKumar SK, Yu L, Taylor CG, Zahradka P and Netticadan T. 2012. Treatment with low-dose resveratrol reverses cardiac impairment in obese prone but not in obese resistant rats. *J Nutr Biochem* 23: 1163-1169
- Marton O, Koltai E, Nyakas C, Bakonyi T, Zenteno-Savin T, Kumagai S, Goto S and Radak Z. 2010. Aging and exercise affect the level of protein acetylation and SIRT1 activity in cerebellum of male rats. *Biogerontology* 11: 679-686
- Merrill GF, Kurth EJ, Hardie DG and Winder WW. 1997. AICA riboside increases AMP-activated protein kinase, fatty acid oxidation, and glucose uptake in rat muscle. *Am J Physiol* 273: E1107-1112
- Nakae J, Cao Y, Hakuno F, Takemori H, Kawano Y, Sekioka R, Abe T, Kiyonari H, Tanaka T, Sakai J, Takahashi S and Itoh H. 2012. Novel repressor regulates insulin sensitivity through interaction with Foxo1. *EMBO J* 31: 2275-2295
- Nasrin N, Wu X, Fortier E, Feng Y, Bare OC, Chen S, Ren X, Wu Z, Streeper RS and Bordone L. 2010. SIRT4 regulates fatty acid oxidation and mitochondrial gene expression in liver and muscle cells. *J Biol Chem* 285: 31995-32002
- Nemoto S, Fergusson MM and Finkel T. 2005. SIRT1 functionally interacts with the metabolic regulator and transcriptional coactivator PGC-1 α . *J Biol Chem* 280: 16456-16460
- Ngo JK and Davies KJ. 2009. Mitochondrial Lon protease is a human stress protein. *Free Radic Biol Med* 46: 1042-1048
- Nunn AV, Guy GW and Bell JD. 2010. Endocannabinoids, FOXO and the metabolic syndrome: redox, function and tipping point—the view from two systems. *Immunobiology* 215: 617-628
- Otera H and Mihara K. 2011. Molecular mechanisms and physiologic functions of mitochondrial dynamics. *J Biochem* 149: 241-251
- Radak Z, Atalay M, Jakus J, Boldogh I, Davies K and Goto S. 2009. Exercise improves import of 8-oxoguanine DNA glycosylase into the mitochondrial matrix of skeletal muscle and enhances the relative activity. *Free Radic Biol Med* 46: 238-243

- Radak Z, Chung HY, Naito H, Takahashi R, Jung KJ, Kim HJ and Goto S. 2004. Age-associated increase in oxidative stress and nuclear factor kappaB activation are attenuated in rat liver by regular exercise. *FASEB J* 18: 749-750
- Radak Z, Zhao Z, Koltai E, Ohno H and Atalay M. 2013. Oxygen Consumption and Usage During Physical Exercise: The Balance Between Oxidative Stress and ROS-Dependent Adaptive Signaling. *Antioxid Redox Signal* 18: 1208-1246.
- Rodgers JT, Lerin C, Gerhart-Hines Z and Puigserver P. 2008. Metabolic adaptations through the PGC-1 alpha and SIRT1 pathways. *FEBS Lett* 582: 46-53
- Romanello V, Guadagnin E, Gomes L, Roder I, Sandri C, Petersen Y, Milan G, Masiero E, Del Piccolo P, Foretz M, Scorrano L, Rudolf R and Sandri M. 2010. Mitochondrial fission and remodelling contributes to muscle atrophy. *EMBO J* 29: 1774-1785
- Rottgers K, Zufall N, Guiard B and Voos W. 2002. The ClpB homolog Hsp78 is required for the efficient degradation of proteins in the mitochondrial matrix. *J Biol Chem* 277: 45829-45837
- Sarkar D, Park ES, Emdad L, Randolph A, Valerie K and Fisher PB. 2005. Defining the domains of human polynucleotide phosphorylase (hPNPaseOLD-35) mediating cellular senescence. *Mol Cell Biol* 25: 7333-7343
- Scheckhuber CQ, Erjavec N, Tinazli A, Hamann A, Nystrom T and Osiewacz HD. 2007. Reducing mitochondrial fission results in increased life span and fitness of two fungal ageing models. *Nat Cell Biol* 9: 99-105
- Smith JJ, Kenney RD, Gagne DJ, Frushour BP, Ladd W, Galonek HL, Israelian K, Song J, Razvadauskaite G, Lynch AV, Carney DP, Johnson RJ, Lavu S, Iffland A, Elliott PJ, Lambert PD, Elliston KO, Jirousek MR, Milne JC and Boss O. 2009. Small molecule activators of SIRT1 replicate signaling pathways triggered by calorie restriction in vivo. *BMC Syst Biol* 3: 31
- Spina RJ, Chi MM, Hopkins MG, Nemeth PM, Lowry OH and Holloszy JO. 1996. Mitochondrial enzymes increase in muscle in response to 7-10 days of cycle exercise. *J Appl Physiol* 80: 2250-2254
- Thyfault JP, Rector RS, Uptergrove GM, Borengasser SJ, Morris EM, Wei Y, Laye MJ, Burant CF, Qi NR, Ridenhour SE, Koch LG, Britton SL and Ibdah JA. 2009. Rats selectively bred for low aerobic capacity have reduced hepatic mitochondrial oxidative capacity and susceptibility to hepatic steatosis and injury. *J Physiol* 587: 1805-1816
- Wang G, Chen HW, Oktay Y, Zhang J, Allen EL, Smith GM, Fan KC, Hong JS, French SW, McCaffery JM, Lightowlers RN, Morse HC, 3rd, Koehler CM and Teitell MA. 2010. PNPASE regulates RNA import into mitochondria. *Cell* 142: 456-467
- Westermann B. 2010. Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol Cell Biol* 11: 872-884
- Wisloff U, Najjar SM, Ellingsen O, Haram PM, Swoap S, Al-Share Q, Fernstrom M, Rezaei K, Lee SJ, Koch LG and Britton SL. 2005. Cardiovascular risk factors emerge after artificial selection for low aerobic capacity. *Science* 307: 418-420
- Yu T, Robotham JL and Yoon Y. 2006. Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology. *Proc Natl Acad Sci U S A* 103: 2653-2658
- Zern TL, Wood RJ, Greene C, West KL, Liu Y, Aggarwal D, Shachter NS and Fernandez ML. 2005. Grape polyphenols exert a cardioprotective effect in pre- and postmenopausal women by lowering plasma lipids and reducing oxidative stress. *J Nutr* 135: 1911-1917
- Zheng J, Chen LL, Zhang HH, Hu X, Kong W and Hu D. 2012. Resveratrol improves insulin resistance of catch-up growth by increasing mitochondrial complexes and antioxidant function in skeletal muscle. *Metabolism* 61: 954-965